

## Abstract

The advent of multiple drug resistant bacterial strains and the lack of novel antibiotic therapeutics to respond remains one of the pressing medical concerns of our time. Recent work has highlighted a class of synthetic compounds, known as antimicrobial lipopeptides, which show great promise as a scaffold for future drug design. One such compound, palmitoyl-Lys-Gly-Gly-DLys (C16-KGGK), has micromolar minimum inhibitory concentrations against a variety of bacterial and fungal species. Here we have used coarse-grained and all-atom molecular dynamics in tandem to probe the biophysical mechanism of action behind this lipopeptide with varying detail and time-scales. Our results are validated to experimental results and suggest a possible mechanism by which lipid bilayers are disrupted.

### **Lipopeptides of Interest**

- Synthetic antimicrobial lipopeptide C16-KGGK
- KGG<mark>K</mark> peptide sequence
- Last lysine is a D-amino acid (in red) - Palmitoyl chain linked via peptide bond
- Physical features bestow selectivity/potency - Net positive charge
- Amphipathic
- Investigated by the Shai lab - Makovitzki, Avrahami, and Shai, PNAS, 2006, 103, 15997-16002

# Multiscale Modeling Approach

- Full atomic detail in a simulation is computationally expensive
- Abstracted detail trades accuracy for sampling
- Multiscale modeling
- Choose the representation that best answers the question
- Multiple resolutions can be effective



#### **All-atom Systems**

- CHARMM27 force field - 2 fs timestep
- 20 lipopeptides - Free in solution above bilayer - Simulates binding
- 180 lipids in bilayer
- 2:1 POPE:POPG (bacteria-like)
- 12,000 water molecules
- 100 mM NaCl
- 60,850 atoms
- 15 ns/day on BG/L - 2048 cores
- 3 simulations - 6 µs total

- **Coarse-grained Systems**
- MARTINI force field - 10 fs timestep
- 48 lipopeptide micelle - Equilibrated in solution
- 480 lipids in bilayer
- 2:1 POPE:POPG (bacteria-like)
- 24,000 water beads
- 100 mM NaCl
- equivalent to 120k atoms
- 100 ns/day on cluster - 8 cores
- 4 simulations - 20 µs total (80 µs effective time)



For reference, use the code to the left to access a digital copy of this poster. To learn more about this work or to contact the authors, visit us at:

# Elucidating Antimicrobial Lipopeptide Action via Combined **Coarse-grained and All-atom Molecular Dynamics** Joshua N. Horn, Jesse D. Sengillo, Alan Grossfield University of Rochester Medical School, Rochester, NY, USA



- One trajectory shown, normalized by area and frames - Left image shows bound leaflet, right shows unbound - 0,0 represents position of the micelle center of mass
- High concentration of POPG directly beneath micelle
- Bilayer reorganizes laterally

• Neat bilayer and unbound leaflets have identical distribution - Diffuse, with very wide peaks

• Drastic change in bound leaflet, sharp peaks at 100° and 180° - Shows ordered headgroup packing

• Image shows a row of lipids ordered by lipopeptide action - Direct interaction between lysines and anionic headgroups



### **Limitations of MARTINI**





• Left shows the micelle above an average 3D density map of the two lipid components, right shows the same with micelle removed

- Green is peptide and red is palmitoyl tail of lipopeptide - Orange and pink are bulk densities of POPE and POPG, respectively - Darker pink in right image indicates regions of higher POPG density • Bilayer crystallization is a cause for concern when working with the MARTINI forcefield

- Consider using the published polarizable model in the future

### **Multiscale Summary**

• C16-KGGK tends to micellize in solution

- Structurally, the lipopeptide resembles a detergent

• All-atom simulations demonstrate that:

- Free lipopeptides will bind and rapidly insert

- Lysines drive initial interactions with the bilayer
- Fully inserted lipopeptides change lipid order parameters • Coarse-grained simulations demonstrate that:

- Lipopeptides spontaneously form large micelles

- Micelles are pseudo-stable and do not readily insert
- Bound micelles demix the bilayer

# **Proposed Model of Action**

• C16-KGGK action is likely a combination of effects

• At high concentrations

- All lipopeptides insert and affect bilayer stability • At lower concentrations

- Available lipopeptides micellize and bind to the bilayer - On longer timescales, micelle demixes lipids

### **Future Plans**

• High concentration all-atom simulations

- Currently in progress

- Quantify C16-KGGK effects on bilayer stability
- Mixed coarse-grained/all-atom sampling

- Take advantage of CG speed to sample binding process - Reconstruct atomistic details at snapshots along trajectory - Run all-atom simulations to gather detailed information - Can parallelize long timescale all-atom simulations • Implement polarizable water into simulations - Expect crystal artifacts to vanish

Some simulation details and results are in press: Horn et al., Biochim Biophys Acta, 2012 Feb; 1818(2):212-8.



Analysis done using LOOS (Lightweight Object Oriented **LIDDS** Structure analysis library), an open source C++ library designed and maintained but he C designed and maintained by the Grossfield lab. LOOS provides a framework for designing analysis tools that interface with file formats of most simulation packages.